

Local correlations in the 1D Bose gas from a scaling limit of the XXZ chain

Balázs Pozsgay

Institute for Theoretical Physics, Universiteit van Amsterdam
Science Park 904, Postbus 94485, 1090 GL Amsterdam, The Netherlands

August 2011

Abstract

We consider the K -body local correlations in the (repulsive) 1D Bose gas for general K , both at finite size and in the thermodynamic limit. Concerning the latter we develop a multiple integral formula which applies for arbitrary states of the system with a smooth distribution of Bethe roots, including the ground state and finite temperature Gibbs-states. In the cases $K \leq 4$ we perform the explicit factorization of the multiple integral. In the case of $K = 3$ we obtain the recent result of Kormos et.al., whereas our formula for $K = 4$ is new. Numerical results are presented as well.

1 Introduction

The delta-function interacting 1D Bose gas (also known as the Lieb-Liniger model or the Quantum Nonlinear Schrödinger equation) is one of the oldest and most important integrable models. Its study goes back to the papers [1, 2] where it was shown that the spectrum can be obtained by the Bethe Ansatz [3]. The thermodynamical properties of the model were determined in [4] using the method nowadays known as the Thermodynamical Bethe Ansatz (TBA). After these seminal papers tremendous effort was devoted to the calculation of correlation functions using various approaches [5, 6, 7, 8, 9, 10, 11]. One of the most important recent results is the exact determination of the long-distance behaviour of correlations [12, 13, 14, 15].

Apart from purely academic interest, the study of the 1D Bose gas was spurred by the recent success of experiments with cold atoms in quasi one-dimensional traps [16, 17, 18, 19, 20, 21, 22]. A remarkable result was presented in [19], where the authors managed to measure exact predictions of the TBA (for further developments and open questions see [23]). In experimental situations the local correlations are of special interest, for example the three-body local correlation is related to the rate of particle loss [24, 25, 22] and to the third moment of the density fluctuations [26, 27]. Moreover, even the four-body correlations might be accessible to experiment, as it was recently demonstrated in a 3D experiment [21].

Concerning the general K -body local correlations (for the precise definition see the main text) there has been considerable theoretical progress, too. The $K = 1$ case is simply given by the (linear) density of particles, whereas the $K = 2$ case was related to the thermodynamical quantities of the model in [28]. Concerning the higher-body cases small-coupling and large-coupling expansions were performed in [28, 29], whereas the exact ground state value of the three-body correlation was calculated in [30]. A new approach was initiated in [31, 32], where an infinite integral series (also called the LeClair-Mussardo or LM series) was derived using a special non-relativistic limit of the sinh-Gordon model. The LM series applies for any K and arbitrary temperature, including the ground state, and it can be considered as an effective large-coupling expansion of the quantity in question. The papers [33, 34] considered the relation between the LM series and previous form factor calculations with the Algebraic

Bethe Ansatz (ABA); in [34] it was shown that the LM series can be understood and proven within the ABA. However, there was one crucial problem: there were no explicit and general results available for the form factors entering the LM series; the numerical results in [31, 32] were obtained using a truncation of the full series.

The important task of the exact summation of the LM series was performed for the first time in the recent article [35], where the authors evaluated the three-body correlation based on a well-supported conjecture for the corresponding form factors. To our best knowledge this is the first time that an exact, explicit and compact result was given for a non-trivial correlation of the 1D Bose gas, valid at arbitrary couplings and temperatures.

In the present work we contribute to the calculation of the K -body correlators using a different approach. Our strategy is the following. First we consider a related physical quantity (the so-called “emptiness formation probability”) on a generic XXZ spin chain and show that a special scaling limit of the spin chain [11, 36] yields the desired correlations in the Bose gas (Section 3). The matrix elements of the operator on the spin chain are calculated in Section 4 borrowing results from the works [37, 38]. We then perform the scaling limit towards the Bose gas in Section 5, this way we obtain the form factors in a finite volume, with a finite number of particles. Finally, the thermodynamic limit is performed in the Bose gas (Section 6) leading to the multiple integral (6.7), which is the main result of this work (see (6.10) for the dimensionless form).

In principle the multiple integrals could be evaluated for any K , but in practice this becomes more and more difficult with increasing K , therefore it is desirable to derive more compact results. In Section 7 we show how to factorize the multiple integral in the cases $K \leq 4$. The results are the expressions (7.3), (7.10) and (7.12). In subsection 7.5 we also present examples of the numerical results.

Finally in Section 8.3 we determine all form factors entering a modified form of the LM series, making it an explicit integral series for the K -body local correlations.

2 The Lieb Liniger model

The second quantized form of the Hamiltonian is

$$H_{LL} = \int_0^L dx \left(\partial_x \Psi^\dagger \partial_x \Psi + c \Psi^\dagger \Psi^\dagger \Psi \Psi \right). \quad (2.1)$$

Here L is the size of the system, periodic boundary conditions are understood and $\Psi(x, t)$ and $\Psi^\dagger(x, t)$ are canonical non-relativistic Bose fields satisfying

$$[\Psi(x, t), \Psi^\dagger(y, t)] = \delta(x - y). \quad (2.2)$$

We used the conventions $m = 1/2$ and $\hbar = 1$ and $c > 0$ is the coupling constant.

The eigenstates of the Hamiltonian (2.1) can be constructed using the Bethe Ansatz [1, 2, 7]. The N -particle coordinate space wave function is given by

$$\chi_N(p|x) = \frac{1}{\sqrt{N!}} \sum_{\mathcal{P}} \exp \left\{ i \sum_j x_j (\mathcal{P}p)_j \right\} \prod_{j>k} \frac{(\mathcal{P}p)_j - (\mathcal{P}p)_k - ic\epsilon(x_j - x_k)}{(\mathcal{P}p)_j - (\mathcal{P}p)_k}, \quad (2.3)$$

where $\epsilon(x)$ is the sign function.

Periodic boundary conditions force the quasi-momenta to be solutions of the Bethe Ansatz equations

$$e^{ip_j L} \prod_{k \neq j} \frac{p_j - p_k - ic}{p_j - p_k + ic} = 1. \quad (2.4)$$

The energy and momentum of the multi-particle state is given by

$$E_N = \sum_j p_j^2 \quad P_N = \sum_j p_j.$$

The norm of the wave function (2.3) is [39, 40]

$$\mathcal{N}^{LL} = \int |\chi_N|^2 = \prod_{j < k} \frac{(p_j - p_k)^2 + c^2}{(p_j - p_k)^2} \times \det \mathcal{G}^{LL} \quad (2.5)$$

with

$$\mathcal{G}_{jk}^{LL} = \delta_{j,k} \left(L + \sum_{l=1}^N \varphi(p_j - p_l) \right) - \varphi(p_j - p_l) \quad (2.6)$$

and

$$\varphi(u) = \frac{2c}{u^2 + c^2}.$$

We will be interested in the matrix elements of the operators

$$\mathcal{O}_K = (\Psi^\dagger(0))^K (\Psi(0))^K.$$

In coordinate space the matrix elements are given by the integrals

$$\langle \phi_N | \mathcal{O}_K | \chi_N \rangle = \frac{N!}{K!(N-K)!} \times \int_0^L dx_1 \dots dx_{N-K} \phi_N^*(0, \dots, 0, x_1, \dots, x_{N-K}) \chi_N(0, \dots, 0, x_1, \dots, x_{N-K}). \quad (2.7)$$

The expectation value of $\mathcal{O}_K$ describes the probability to have K particles at the same point. It is useful to introduce the dimensionless quantities

$$g_K = \frac{\langle \mathcal{O}_K \rangle}{n^K},$$

where $n = N/L$ is the particle density. It can be shown by scaling arguments that in the thermodynamic limit g_K only depends on the dimensionless parameters

$$\gamma = \frac{c}{n} \quad \tau = \frac{T}{n^2},$$

where T is the temperature (we used the convention $k_B = 1$ for the Boltzmann constant).

In principle the form factors (2.7) could be obtained by performing the integrals in coordinate space, but this becomes increasingly complicated with growing N . Note also that the Algebraic Bethe Ansatz for the Bose gas does not lead to simple results either: the action of the field operators on Bethe states can be evaluated easily, but afterwards one would have to compute scalar products of Bethe states with a reduced set of $N - K$ particles, neither of which are on-shell, and there is no good formula for the scalar products of such states. One way out of these problems is to consider a related quantity (the “emptiness formation probability”) on the XXZ spin chain, where there are methods available to compute its matrix elements.

3 The XXZ chain and its special scaling limit

The XXZ spin chain with M sites and periodic boundary conditions is given by the following Hamiltonian:

$$H = J \sum_{j=1}^M (S_j^x S_{j+1}^x + S_j^y S_{j+1}^y + \Delta (S_j^z S_{j+1}^z - 1/4)) + h \sum_{j=1}^M S_j^z. \quad (3.1)$$

This model is also solvable by the Bethe Ansatz [3, 41, 42, 43]. The N -particle eigenstates are given by

$$|\phi_N\rangle = \frac{1}{\sqrt{N!}} \sum_{y_1=1}^L \dots \sum_{y_N=1}^L \phi_N(\lambda | y_1, \dots, y_N) \sigma_{y_1}^- \dots \sigma_{y_N}^- |0\rangle. \quad (3.2)$$

Here $|0\rangle$ is the reference state with all spins up and y_j are the positions of the down spins. The amplitudes are

$$\phi_N(\lambda|y) = \frac{1}{\sqrt{N!}} \sum_{\mathcal{P} \in \sigma_N} \prod_{1 \leq m < n \leq N} \frac{\sinh((\mathcal{P}\lambda)_m - (\mathcal{P}\lambda)_n + \epsilon(y_n - y_m)\eta)}{\sinh((\mathcal{P}\lambda)_m - (\mathcal{P}\lambda)_n)} \prod_{l=1}^N F((\mathcal{P}\lambda)_l, y_l), \quad (3.3)$$

where

$$F(\lambda, y) = \frac{1}{\sinh(\lambda - \xi_y)} \prod_{j=1}^{y-1} \frac{\sinh(\lambda - \xi_j + \eta)}{\sinh(\lambda - \xi_j)}. \quad (3.4)$$

The parameter η is related to the anisotropy:

$$\Delta = \cosh \eta.$$

In (3.4) we introduced inhomogeneities ξ_j for the sites of the spin chain; they will be used as a technical tool to obtain the form factors in section 4. The physical limit consists of setting all $\xi_j \rightarrow \eta/2$. The expression (3.3) is a seemingly over-complicated way to write down the wave function, because it is valid at arbitrary values of the variables y_j and not only in the region $y_1 < \dots < y_M$. We used this form to have an exact agreement with the conventions used in (2.3).

The Bethe equations follow from the periodicity of the wave function and they read

$$d(\lambda_j) \prod_{k \neq j} \frac{\sinh(\lambda_j - \lambda_k + \eta)}{\sinh(\lambda_j - \lambda_k - \eta)} = 1, \quad (3.5)$$

where

$$d(\lambda) = \prod_{k=1}^M \frac{\sinh(\lambda - \xi_k)}{\sinh(\lambda - \xi_k + \eta)}. \quad (3.6)$$

In the normalization (3.2)-(3.3) the norm of the wave function is given by

$$\begin{aligned} \mathcal{N}^{XXZ} &= \sum_{y_1} \dots \sum_{y_N} |\phi(y_1, \dots, y_N)|^2 = \\ &= (-\sinh \eta)^{-N} \prod_{j < k} f(\lambda_j, \lambda_k) f(\lambda_k, \lambda_j) \times \det \mathcal{G}^{XXZ} \end{aligned} \quad (3.7)$$

with

$$\mathcal{G}_{jk}^{XXZ} = \delta_{j,k} \left(\frac{d'(\lambda_j)}{d(\lambda_j)} + \sum_{l \neq j} \varphi^{XXZ}(\lambda_j - \lambda_l) \right) - \varphi^{XXZ}(\lambda_j - \lambda_k). \quad (3.8)$$

The kernel φ^{XXZ} is given by

$$\varphi^{XXZ}(u) = \frac{-\sinh \eta}{\sinh(u + \eta/2) \sinh(u - \eta/2)}. \quad (3.9)$$

One-particle momenta and energies are given by the formulas

$$e^{ip(\lambda)} = \frac{\sinh(\lambda + \eta/2)}{\sinh(\lambda - \eta/2)} \quad e(\lambda) = J \frac{\sinh^2 \eta}{\cos(2\lambda) - \cosh \eta} - h.$$

3.1 Towards the Lieb-Liniger model

There is a special scaling limit of the XXZ chain which yields the physical quantities of the Lieb-Liniger model [44, 36, 11]. In order to obtain the Bose gas in a finite volume L one has to set

$$\eta = i\pi - i\varepsilon \quad M = \frac{c}{\varepsilon^2} L$$

and let $\varepsilon \rightarrow 0$ (here c is the coupling constant of the Bose gas). The number N of the magnons has to be kept fixed and the rapidities of the particles have to be scaled as

$$\lambda_j = p_j \frac{\varepsilon}{c}.$$

After the limiting procedure the magnons can be identified as the particles of the Bose gas with rapidity p_j . It can be shown that under an appropriate scaling of the parameters J and h

$$e(\lambda) = J \frac{\sinh^2 \eta}{\cos(2\lambda) - \cosh \eta} - h \quad \rightarrow \quad p^2 - \mu,$$

where μ is the chemical potential in the Bose gas. However, this will be not needed in the following; we will consider the Bethe wave functions and the form factors of local operators. In the following we assume that the homogeneous limit $\xi_j \rightarrow \eta/2$ is performed first on the spin chain, and the limit towards the Bose gas is taken afterwards.

Taking the scaling limit of the Bethe equations (3.5) results in

$$(-1)^M e^{i\nu_j l} \prod_{k \neq j} \frac{\nu_j - \nu_k - ic}{\nu_j - \nu_k + ic} = 1. \quad (3.10)$$

For the sake of simplicity we only consider even chains so that no twist appears in the Bethe equations.

The limiting form of the Bethe wave function can be taken by setting $x_j = \frac{\varepsilon^2}{c} y_j$ and keeping x_j finite, which will correspond to the position of the particles of the Bose gas. The wave function then reads

$$\Psi_N(x|\nu) = \frac{1}{\sqrt{N!}} \sum_{\mathcal{P} \in \sigma_N} \prod_{m > n} - \frac{(\mathcal{P}\nu)_m - (\mathcal{P}\nu)_n + i\varepsilon(x_m - x_n)c}{(\mathcal{P}\nu)_m - (\mathcal{P}\nu)_n} \prod_{l=1}^N F((P\nu)_l, x_l), \quad (3.11)$$

where

$$F(\nu, x) = e^{-i\nu x} (-1)^y. \quad (3.12)$$

Apart from factors of (-1) the above expression is equal to the complex conjugate of the Bethe wave function (2.3). It can be argued that the factors of $(-1)^{y_j}$ don't affect the calculation of form factors of local operators. Indeed, for any coordinate space calculation one has to take the product of two wave functions with the down spins placed at prescribed positions. Depending on the operator in question an overall factor of (-1) may remain, but the position dependent factors of $(-1)^{y_j}$ always cancel. For the operators considered in this paper every such factor cancels, therefore they will be neglected in the following.

Due to the relation between y and x it is expected that the norm of the wave function behaves as

$$\mathcal{N}^{XXZ} \rightarrow \left(\frac{c}{\varepsilon^2}\right)^N \mathcal{N}^{LL}. \quad (3.13)$$

Comparing the formulas (2.5) and (3.7) we obtain the same scaling: the Gaudin determinants behave as

$$\det \mathcal{G}^{XXZ} \rightarrow \left(\frac{ic}{\varepsilon}\right)^N \det \mathcal{G}^{LL}$$

and the prefactors contribute an extra $(i\varepsilon)^{-N}$.

3.2 The emptiness formation probability

We are interested in the local operators $E_j^{\alpha\beta}$ acting on site j with matrix elements

$$(E^{\alpha\beta})_{kl} = \delta_{k,\alpha} \delta_{l,\beta}.$$

In particular we consider the composite operator

$$s_K = E_1^{--} E_2^{--} \dots E_K^{--}. \quad (3.14)$$

When sandwiched between two states, this operator forces K particles to occupy the first K sites. The expectation value of s_K (or sometimes its spin reverse) is called the “emptiness formation probability”.

We will show that the operator s_K scales to $\mathcal{O}_K$ in the limiting procedure. In the coordinate Bethe Ansatz its N -particle form factors are given by

$$\langle \{\lambda\} | s_K | \{\mu\} \rangle = \frac{N!}{K!(N-K)!} \times \sum_{y_1, \dots, y_{N-K}=K+1}^M \phi_N^*(\lambda | 1, 2, \dots, K, y_1, \dots, y_{N-K}) \phi_N(\mu | 1, 2, \dots, K, y_1, \dots, y_{N-K}). \quad (3.15)$$

This formula has to be compared to (2.7). It is easy to see that the scaling limit of the Bethe wave functions works even if a fixed number of particles are placed on the first few sites:

$$\phi_N(\mu | 1, 2, \dots, K, y_1, \dots, y_{N-K}) \rightarrow \chi_N(p | 0, 0, \dots, 0, x_1, \dots, x_{N-K})^*.$$

Therefore the un-normalized form factor will behave as

$$\left(\frac{\varepsilon^2}{c}\right)^{M-K} \langle \{\lambda\} | s_K | \{\mu\} \rangle \rightarrow \langle \{p\} | \mathcal{O}_K | \{k\} \rangle^*,$$

where it is understood that

$$\frac{c}{\varepsilon} \lambda_j \rightarrow p_j \quad \frac{c}{\varepsilon} \mu_j \rightarrow k_j.$$

For the normalized form factors we get

$$\left(\frac{\varepsilon^2}{c}\right)^{-K} \frac{\langle \{\lambda\} | s_K | \{\mu\} \rangle}{\sqrt{\langle \lambda | \lambda \rangle \langle \mu | \mu \rangle}} \rightarrow \frac{\langle \{p\} | \mathcal{O}_K | \{k\} \rangle^*}{\sqrt{\langle p | p \rangle \langle k | k \rangle}}. \quad (3.16)$$

Our strategy is to obtain explicit determinant formulas for the matrix elements (3.15) and to take the scaling limit according to (3.16).

4 Form factors in the XXZ chain

In this section we compute explicit determinant formulas for the matrix elements (3.15) in the framework of Algebraic Bethe Ansatz (ABA). Mostly we will use the results of the papers [37, 38]; the only difference between the present approach and the traditional methods is that here the homogeneous limit $\xi_j \rightarrow \eta/2$ is taken explicitly before performing the thermodynamic limit or the limit towards the Bose gas.

The central object in ABA is the monodromy matrix, a 2×2 matrix in the so-called auxiliary space with operator valued entries which act on the Hilbert space of the spin chain:

$$T(u) = \begin{pmatrix} A(u) & B(u) \\ C(u) & D(u) \end{pmatrix}.$$

It is built from the so-called local L-matrices:

$$T(u) = L_M(u) \dots L_1(u),$$

where

$$L_j(u) = R_{0j}(u - \xi_j).$$

Here j refers to the quantum space of the spin at site j and 0 refers to the auxiliary space and the parameters ξ_j are identical to the inhomogeneities already introduced in (3.4). The operator $R(u)$ is the R-matrix of the XXZ type:

$$R(u) = \frac{1}{\sinh(u + \eta)} \begin{pmatrix} \sinh(u + \eta) & & & \\ & \sinh(u) & \sinh(\eta) & \\ & \sinh(\eta) & \sinh(u) & \\ & & & \sinh(u + \eta) \end{pmatrix}. \quad (4.1)$$

The trace of the monodromy matrix is called the transfer matrix:

$$\tau(u) = A(u) + D(u).$$

In the homogeneous limit $\xi_j = \eta/2$ it is related to the Hamiltonian (3.1) at $h = 0$ as

$$H \sim \left. \frac{d}{du} \tau(u) \right|_{u=\eta/2} + \text{const.}$$

The normalization (4.1) of the R -matrix results in the following vacuum eigenvalues:

$$A(u)|0\rangle = |0\rangle \quad D(u)|0\rangle = d(u)|0\rangle,$$

with $d(\lambda)$ given by (3.6).

In the framework of ABA the Bethe states are

$$\langle 0 | \prod_{j=1}^N C(\lambda_j) \quad \text{and} \quad \prod_{j=1}^N B(\mu_j) | 0 \rangle.$$

They are eigenstates of the transfer matrix if the rapidities satisfy the Bethe equations (3.5). Apart from an overall normalization factor they are identical to the states given by the coordinate wave functions (3.2).

In order to compute the matrix elements of s_K in the framework of ABA the local operators have to be expressed in terms of the entries of the transfer matrix. This problem was solved in [38, 45] leading to the following theorem:

$$E_j^{\alpha\beta} = \prod_{k=1}^{j-1} (A + D)(\xi_k) \times T^{\alpha\beta}(\xi_j) \times \prod_{k=j+1}^M (A + D)(\xi_k). \quad (4.2)$$

Applying this formula to operators E_j^{--} on neighbouring sites one gets

$$s_k = D(\xi_1)D(\xi_2) \dots D(\xi_k) \prod_{l=k+1}^L (A + D)(\xi_l). \quad (4.3)$$

Evaluated on Bethe states equation (4.3) yields

$$\langle 0 | \prod_{j=1}^N C(\lambda_j) s_k \prod_{j=1}^N B(\mu_j) | 0 \rangle = \langle 0 | \prod_{j=1}^N C(\lambda_j) D(\xi_1)D(\xi_2) \dots D(\xi_k) \prod_{j=1}^N B(\mu_j) | 0 \rangle \times \prod_{j=1}^k \frac{1}{t(\xi_j, \{\mu\})} \quad (4.4)$$

with $t(u, \{\mu\})$ being the corresponding eigenvalue of the transfer matrix. Evaluated at the inhomogeneities it reads

$$t(\xi_j, \{\mu\}) = \prod_{k=1}^N \frac{\sinh(\mu_k - \xi_j + \eta)}{\sinh(\mu_k - \xi_j)}. \quad (4.5)$$

In (4.4) we also used the fact that

$$\prod_{l=1}^L (A + D)(\xi_l) = 1.$$

The action of multiple D operators on the dual state results in [37]

$$\begin{aligned} \langle 0 | \prod_{j=1}^N C(\lambda_j) D(\xi_1)D(\xi_2) \dots D(\xi_k) = \\ \sum_{\substack{\{\lambda^+\} \cup \{\lambda^-\} \\ |\{\lambda^+\}|=k}} \frac{\prod_{o,p} \sinh(\lambda_p^+ - \xi_o + \eta)}{\prod_{j < l} \sinh(\xi_l - \xi_j) \sinh(\lambda_j^+ - \lambda_l^+)} \times \det t(\lambda_l^+, \xi_j) \times \prod_{o,p} f(\lambda_p^+, \lambda_o^-) \times \\ \prod_l d(\lambda_l^+) \langle 0 | \prod_o C(\lambda_o^-) C(\xi_1) \dots C(\xi_k) \end{aligned}$$

with

$$t(\lambda, \xi) = \frac{\sinh \eta}{\sinh(\lambda - \xi) \sinh(\lambda - \xi + \eta)}.$$

The scalar product of an arbitrary state and a Bethe state is [46]

$$\langle 0 | \prod_j C(\lambda_j) \prod_j B(\mu_j) | 0 \rangle = \frac{\prod_{j,k} \sinh(\mu_j - \lambda_k + \eta)}{\prod_{j < k} \sinh(\lambda_j - \lambda_k) \sinh(\mu_k - \mu_j)} \times \det S, \quad (4.6)$$

where

$$S_{jk} = t(\mu_j, \lambda_k) - d(\lambda_k) t(\lambda_k, \mu_j) \prod_{l=1}^N \frac{\sinh(\lambda_k - \mu_l + \eta)}{\sinh(\lambda_k - \mu_l - \eta)}.$$

Specializing this to the present case

$$\begin{aligned} \langle 0 | \prod_o C(\lambda_o^-) C(\xi_1) \dots C(\xi_k) \prod_j B(\mu_j) | 0 \rangle = \\ \frac{\prod_{j,k} \sinh(\mu_j - \lambda_k^- + \eta) \prod_{j,o} \sinh(\mu_j - \xi_o + \eta)}{\prod_{j < k} \sinh(\mu_k - \mu_j) \prod_{j < k} \sinh(\lambda_j^- - \lambda_k^-) \prod_{j < k} \sinh(\xi_j - \xi_k) \prod_{j,k} \sinh(\xi_j - \lambda_k)} \times \\ \times \det U. \end{aligned}$$

Here

$$\begin{aligned} U_{jl} = t(\mu_j, \xi_l) \quad \text{for } l = 1 \dots K \\ U_{j,k+l} = t(\mu_j, \lambda_l^-) - d(\lambda_l^-) t(\lambda_l^-, \mu_j) \prod_{o=1}^N \frac{\sinh(\lambda_l^- - \mu_o + \eta)}{\sinh(\lambda_l^- - \mu_o - \eta)} \quad \text{otherwise.} \end{aligned} \quad (4.7)$$

Therefore, the form factor of the inhomogeneous chain is given by

$$\begin{aligned} \langle \{\lambda\} | s_k | \{\mu\} \rangle = \\ \prod_{j=1}^k \frac{1}{t(\xi_j)} \times \sum_{\substack{\{\lambda^+\} \cup \{\lambda^-\} \\ |\{\lambda^+\}| = k}} \frac{\prod_{o,p} \sinh(\lambda_p^+ - \xi_o + \eta)}{\prod_{j < l} \sinh(\xi_l - \xi_j) \sinh(\lambda_j^+ - \lambda_l^+)} \prod_{o,p} f(\lambda_p^+, \lambda_o^-) \times \prod_l d(\lambda_l^+) \\ \frac{\prod_{j,k} \sinh(\mu_j - \lambda_k^- + \eta) \prod_{j,o} \sinh(\mu_j - \xi_o + \eta)}{\prod_{j < k} \sinh(\mu_k - \mu_j) \prod_{j < k} \sinh(\lambda_j^- - \lambda_k^-) \prod_{j < k} \sinh(\xi_j - \xi_k) \prod_{j,k} \sinh(\xi_j - \lambda_k)} \times \\ \times \det t(\lambda_l^+, \xi_j) \times \det U. \end{aligned} \quad (4.8)$$

We now perform the homogeneous limit $\xi_j \rightarrow \eta/2$ following the method of [47]. For the matrix M we get

$$\lim \frac{\det M}{\prod_{j > l} \sinh(\xi_j - \xi_k)} = \frac{1}{\prod_{\alpha=1}^{m-1} \alpha!} \det H$$

with

$$H_{jl} = \left[\left(\frac{\partial}{\partial \xi} \right)^{l-1} t(\lambda_k, \xi) \right]_{\xi=\eta/2}.$$

It is advantageous to use the form

$$t(\lambda, \xi) = \frac{\cosh(\lambda - \xi)}{\sinh(\lambda - \xi)} - \frac{\cosh(\lambda - \xi + \eta)}{\sinh(\lambda - \xi + \eta)}.$$

Taking the derivatives one is free to replace [48]

$$\left(\frac{\partial}{\partial \xi} \right)^{l-1} t(\lambda, \xi) \rightarrow (-1)^{l-1} (l-1)! \left[\left(\frac{\cosh(\lambda - \xi)}{\sinh(\lambda - \xi)} \right)^l - \left(\frac{\cosh(\lambda - \xi + \eta)}{\sinh(\lambda - \xi + \eta)} \right)^l \right].$$

The same steps can be performed for the corresponding elements of the matrix U . Finally the homogeneous limit reads

$$\begin{aligned} \langle \{\lambda\} | s_K | \{\mu\} \rangle &= \sum_{\substack{\{\lambda^+\} \cup \{\lambda^-\} \\ |\{\lambda^+\}| = K}} \frac{\prod_{o,p} \sinh(\lambda_p^+ + \eta/2)}{\prod_{j < l} \sinh(\lambda_j^+ - \lambda_l^+)} \prod_{o,p} f(\lambda_p^+, \lambda_o^-) \times \prod_l d(\lambda_l^+) \times \\ &\frac{\prod_{j,k} \sinh(\mu_j - \lambda_k^- + \eta) \prod_j \sinh^K(\mu_j - \eta/2)}{\prod_{j < k} \sinh(\mu_j - \mu_k) \prod_{j < k} \sinh(\lambda_j^- - \lambda_k^-) \prod_l \sinh^K(\eta/2 - \lambda_l)} \times \det O \det V. \end{aligned} \quad (4.9)$$

with

$$O_{jl} = \left[\left(\frac{\cosh(\lambda_j^+ - \eta/2)}{\sinh(\lambda_j^+ - \eta/2)} \right)^l - \left(\frac{\cosh(\lambda_j^+ + \eta/2)}{\sinh(\lambda_j^+ + \eta/2)} \right)^l \right] \quad (4.10)$$

and

$$\begin{aligned} V_{jl} &= \left[\left(\frac{\cosh(\mu_j - \eta/2)}{\sinh(\mu_j - \eta/2)} \right)^l - \left(\frac{\cosh(\mu_j + \eta/2)}{\sinh(\mu_j + \eta/2)} \right)^l \right] \quad \text{for } l = 1 \dots K \\ V_{j,K+l} &= t(\mu_j, \lambda_l^-) - d(\lambda_l^-) t(\lambda_l^-, \mu_j) \prod_{o=1}^N \frac{\sinh(\lambda_l^- - \mu_o + \eta)}{\sinh(\lambda_l^- - \mu_o - \eta)} \quad \text{otherwise.} \end{aligned} \quad (4.11)$$

5 The scaling limit of the form factors

Here we take the scaling limit of the formula (4.9) to obtain the matrix elements of $\mathcal{O}_K$ in the Bose gas. We substitute

$$\eta = i\pi - i\varepsilon \quad \lambda = \frac{\varepsilon}{c} p \quad \mu = \frac{\varepsilon}{c} k.$$

It is straightforward to calculate the limiting values of the prefactors, but the determinants need special care. The elements of O read

$$O_{jl} = \left[\left(\frac{\sinh(\lambda_j^+ + i\varepsilon/2)}{\cosh(\lambda_j^+ + i\varepsilon/2)} \right)^l - \left(\frac{\sinh(\lambda_j^+ - i\varepsilon/2)}{\cosh(\lambda_j^+ - i\varepsilon/2)} \right)^l \right]. \quad (5.1)$$

The leading terms will be

$$O_{jl} \rightarrow l \left(\frac{\varepsilon}{c} \right)^{l-1} (p_j^+)^{l-1} i\varepsilon,$$

which yields

$$\det O \rightarrow K! (i\varepsilon)^K \left(\frac{\varepsilon}{c} \right)^{(K-1)K/2} \det [(p_j^+)^{l-1}] = K! (i\varepsilon)^K \left(\frac{\varepsilon}{c} \right)^{(k-1)k/2} \prod_{j>l} (p_j^+ - p_l^+).$$

One can use the same expansion for the first K columns of the matrix V .

To obtain the proper normalization note that in ABA the norm of the Bethe state scales as

$$\langle 0 | \prod_j C(\lambda_j) \prod_j B(\lambda_j) | 0 \rangle \rightarrow c^N \frac{\prod_{j<l} ((p_j - p_l)^2 + c^2)}{\prod_{j<l} (p_j - p_l)^2} \times \det \mathcal{G}_{LL}.$$

This differs from (2.5) by the overall factor of c^N .

Collecting all the factors and performing a complex conjugation we find

$$\begin{aligned} \langle \{p\} | \mathcal{O}_K | \{k\} \rangle &= c^{K-N} (K!)^2 \sum_{\substack{\{p^+\} \cup \{p^-\} \\ |\{p^+\}| = K}} \prod_{o,l} f(p_l^+, p_o^-) \prod_l e^{-iL p_l^+} \times \\ &\times \frac{\prod_{j,l} (k_j - p_l^- + ic)}{\prod_{j<l} (k_j - k_l) \prod_{j<k} (p_j^- - p_k^-)} \times \det Z, \end{aligned} \quad (5.2)$$

with

$$\begin{aligned} Z_{jl} &= (k_j)^{l-1} \quad \text{for } l = 1 \dots K \\ Z_{j,K+l} &= t(k_j, p_l^-) - e^{-iLp_l^-} t(p_l^-, k_j) \prod_{o=1}^N \frac{(p_l^- - k_o + ic)}{(p_l^- - k_o - ic)} \quad \text{otherwise.} \end{aligned} \quad (5.3)$$

Here we used

$$t(u) = \frac{ic}{u(u+ic)}.$$

Equation (5.2) refers to the normalization where the norms of the states $|\{p\}\rangle$ and $|\{k\}\rangle$ are given by (2.5).

The result (5.2) is valid whenever the set $\{k\}$ satisfies the Bethe equations. In the case when $\{p\}$ is also a solution but different from $\{k\}$ we obtain the form factors

$$\begin{aligned} F_N^K(\{p\}, \{k\}) &= c^{K-N} (K!)^2 \sum_{\substack{\{p^+\} \cup \{p^-\} \\ |\{p^+\}|=K}} \prod_{o,l} f(p_o^-, p_l^+) \prod_l \times \\ &\times \frac{\prod_{j,l} (k_j - p_l^- + ic)}{\prod_{j<l} (k_j - k_l) \prod_{j<k} (p_j^- - p_k^-)} \times \det V, \end{aligned} \quad (5.4)$$

with

$$\begin{aligned} V_{jl} &= (k_j)^{l-1} \quad \text{for } l = 1 \dots K \\ V_{j,K+l} &= t(k_j, p_l^-) + t(p_l^-, k_j) \prod_{o=1}^N \frac{(p_l^- - k_o + ic)(p_l^- - p_o - ic)}{(p_l^- - k_o - ic)(p_l^- - p_o + ic)} \quad \text{otherwise.} \end{aligned} \quad (5.5)$$

This is a new result of the present work. In the case of $K = 1$ eq. (5.4) yields an alternative representation for the form factors of the density operator, which were previously determined in [46, 8, 49].

To obtain the mean value of $\mathcal{O}_K$ we take the limit $\{p\} \rightarrow \{k\}$ in (5.2) and divide by the norm (2.5) resulting in

$$\langle \mathcal{O}_K \rangle_N = (K!)^2 \sum_{\substack{\{p^+\} \cup \{p^-\} \\ |\{p^+\}|=K}} \left[\prod_{j>l} \frac{p_j^+ - p_l^+}{(p_j^+ - p_l^+)^2 + c^2} \right] \times \frac{\det \mathcal{H}}{\det \mathcal{G}^{LL}}. \quad (5.6)$$

The elements of $\mathcal{H}$ are given by

$$\mathcal{H}_{j,l} = \begin{cases} (p_j)^{l-1} & \text{for } l = 1 \dots K \\ \mathcal{G}_{j,l}^{LL} & \text{for } l = K+1 \dots N. \end{cases} \quad (5.7)$$

Here it is understood that in both $\mathcal{G}^{LL}$ and $\mathcal{H}$ the ordering of the rapidities is given by $\{p\} = \{\{p^+\}, \{p^-\}\}$. The matrix $\mathcal{H}$ differs from $\mathcal{G}^{LL}$ only in those columns which belong to the subset $\{p^+\}$.

In the case of $K = 1$ the above formula results in

$$\langle \mathcal{O}_1 \rangle = \frac{N}{L}$$

as it should. To prove this note that the sums of the columns of $\mathcal{G}^{LL}$ are equal to L in every row, therefore every $\mathcal{H}$ gives

$$\mathcal{H} = \frac{1}{L} \mathcal{G}^{LL}.$$

6 Expectation values in the thermodynamic limit

In this section we evaluate the thermodynamic limit of (5.6). We consider a Bethe state $|\Omega\rangle$ in a large volume L with a large number of particles such that the particle density $n = N/L$ is fixed. In the thermodynamic limit one defines the density of roots $\rho^{(r)}(p)$ and holes $\rho^{(h)}(p)$ and the total density $\rho(p) = \rho^{(r)}(p) + \rho^{(h)}(p)$. This latter function satisfies the Lieb-equation

$$\rho(p) = \frac{1}{2\pi} + \int_{-\infty}^{\infty} dq \varphi(p-q)f(q)\rho(q). \quad (6.1)$$

Here

$$f(p) = \frac{\rho^{(r)}(p)}{\rho(p)} \quad (6.2)$$

is a distribution function characterizing the state in question. In thermal equilibrium $f(p) = (1 + e^{\varepsilon(p)})^{-1}$ where $\varepsilon(p)$ is the so-called pseudo-energy, which is a solution of the TBA equation

$$\varepsilon(p) = \frac{p^2 - \mu}{T} - \int_{-\infty}^{\infty} \frac{dp'}{2\pi} \varphi(p-p') \log(1 + e^{-\varepsilon(p')}). \quad (6.3)$$

At zero temperature we recover the ground-state distribution

$$f(p) = \begin{cases} 1 & |p| \leq \Lambda \\ 0 & |p| > \Lambda \end{cases} \quad (6.4)$$

with Λ being the Fermi-rapidity. The particle number is always given by the formula

$$n = \frac{N}{L} = \int f(p)\rho(p).$$

We proceed to calculate the thermodynamic limit of (5.6) using the techniques of [50]. The ratio of determinants is calculated as

$$\frac{\det \mathcal{H}}{\det \mathcal{G}^{LL}} = \det \left((\mathcal{G}^{LL})^{-1} \mathcal{H} \right).$$

The resulting matrix on the r.h.s. will be equal to the identity matrix except for those columns belonging to the set $\{p^+\}$. These elements can be evaluated by transforming the action of $\mathcal{G}^{LL}$ into an integral equation. This results in

$$\frac{\det \mathcal{H}}{\det \mathcal{G}^{LL}} = \prod_o \frac{1}{2\pi L \rho(p_o^+)} \times \det I,$$

where

$$I_{jl} = h^{(l-1)}(p_j^+).$$

Here $h^{(l)}(u)$ is the solution of the linear integral equation

$$h^{(l)}(p) = p^l + \int_{-\infty}^{\infty} \frac{dq}{2\pi} \varphi(p-q)f(q)h^{(l)}(q). \quad (6.5)$$

Note that in this normalization $h^{(0)}(p) = 2\pi\rho(p)$ and $f(p)h^{(0)}(p) = 2\pi\rho^{(r)}(p)$.

As a final step one integrates over the rapidities p_j^+ and in the thermodynamic limit one gets

$$\langle \Omega | \mathcal{O}_K | \Omega \rangle = K! \int \frac{dp_1}{2\pi} \dots \frac{dp_K}{2\pi} \prod_o f(p_o) \prod_{j>l} \frac{p_j - p_l}{(p_j - p_l)^2 + c^2} \times \det I. \quad (6.6)$$

The prefactors are completely anti-symmetric therefore one can expand the determinant and write

$$\langle \Omega | \mathcal{O}_K | \Omega \rangle = (K!)^2 \int \frac{dp_1}{2\pi} \dots \frac{dp_K}{2\pi} \prod_{j>l} \frac{p_j - p_l}{(p_j - p_l)^2 + c^2} \prod_{o=1}^K f(p_o) h^{(o-1)}(p_o). \quad (6.7)$$

This formula is the main result of our paper. Its explicit factorization is performed in the cases $K = 2, 3, 4$ in section 7.

For practical purposes it is useful to derive the dimensionless multiple integral for the quantity

$$g_K = \frac{\langle \mathcal{O}_K \rangle}{n^K},$$

which only depends on the dimensionless coupling constant $\gamma = c/n$ and the dimensionless version of the distribution functions $f(p)$. We define

$$q = \frac{p}{c} \quad f(q) = f(p = qc).$$

In the finite temperature case $f(q) = (1 + e^{\tilde{\varepsilon}(q)})^{-1}$ where $\tilde{\varepsilon}(q)$ is the solution of the dimensionless equation

$$\tilde{\varepsilon}(q) = -\alpha + \frac{q^2 \gamma^2}{\tau} - \int_{-\infty}^{\infty} \frac{dq'}{2\pi} \frac{2}{(q - q')^2 + 1} \log(1 + e^{-\varepsilon(p')}), \quad (6.8)$$

with

$$\alpha = \frac{\mu}{T} \quad \tau = \frac{T}{n^2}.$$

Defining the dimensionless functions $\tilde{h}^{(l)}(q)$ as

$$\tilde{h}^{(l)}(q) = q^l + \int_{-\infty}^{\infty} \frac{dq'}{2\pi} \frac{2}{(q' - q)^2 + 1} f(q') \tilde{h}^{(l)}(q') \quad (6.9)$$

we find $\tilde{h}^{(l)}(q) = c^l h^{(l)}(p = qc)$. Thus the dimensionless multiple integral formula is expressed as

$$g_K = (K!)^2 \gamma^K \int \frac{dq_1}{2\pi} \cdots \frac{dq_K}{2\pi} \prod_{j>l} \frac{q_j - q_l}{(q_j - q_l)^2 + 1} \prod_{j=1}^K f(q_j) \tilde{h}^{(j-1)}(q_j). \quad (6.10)$$

6.1 The $c \rightarrow 0^+$ limit

We take the small-coupling limit of the dimensionful formula (6.6) by sending $c \rightarrow 0$ and keeping n fixed. The limiting form of the kernel $\varphi(u)$ is given by

$$\varphi(u) \rightarrow 2\pi \delta(u). \quad (6.11)$$

Therefore the solution of the integral equations (6.5) is

$$h^{(l)}(p) = \frac{p^l}{1 - f(p)}.$$

The determinant in (6.6) has the limiting value

$$\det I \rightarrow \prod_j \frac{1}{1 - f(p)} \prod_{j>k} (p_j - p_k)$$

resulting in

$$\langle \mathcal{O}_K \rangle \rightarrow K! \prod_{j=1}^K \left[\int \frac{dp_j}{2\pi} \frac{f(p_j)}{1 - f(p_j)} \right].$$

Note that

$$\langle \mathcal{O}_1 \rangle = n = \int \frac{dp}{2\pi} \frac{f(p)}{1 - f(p)},$$

therefore

$$g_K = \frac{\langle \mathcal{O}_K \rangle}{n^K} \rightarrow K!$$

as it should be for free bosons.

The above calculation only applies if $f(p) < 1$, therefore the result is not valid for the ground state. In fact

$$\lim_{c \rightarrow 0} \lim_{T \rightarrow 0} g_K = 1.$$

We have checked this property numerically for $K \leq 4$ (see subsection 7.5). To prove it analytically from (6.6) one has to carefully analyze the sub-leading corrections to the integral equation (6.5) with the weight function (6.4). We leave this problem for further research.

6.2 The $c \rightarrow \infty$ limit

Here we derive the leading term in the large coupling expansion of g_K . In the $c \rightarrow \infty$ limit the kernel $\varphi(p)$ is of order $1/c$, therefore to leading order

$$h^{(l)}(p) = p^l$$

and

$$\langle \mathcal{O}_K \rangle \rightarrow \frac{K!}{c^{K(K-1)}} \int \frac{dp_1}{2\pi} \dots \frac{dp_K}{2\pi} \prod_o f(p_o) \prod_{j>l} (p_j - p_l)^2.$$

Evaluating this formula for the ground state gives

$$g_K = \frac{K!}{2^K} \left(\frac{\pi}{\gamma} \right)^{K(K-1)} \times \int_{-1}^1 dx_1 \dots dx_K \prod_{j>l} (x_j - x_l)^2.$$

This result was already obtained in the papers [28, 29, 31, 32].

7 Factorization of the multiple integrals

In this section we perform the factorization of the multiple integral formula (6.7) in the cases $K = 2, 3, 4$ ¹. In order to keep the formulas as short as possible we will use the following notation:

$$\int d\tilde{p} \dots = \int \frac{dp}{2\pi} f(p) \dots$$

Moreover we will suppress the dependence of the mean value on the state $|\Omega\rangle$ and we write simply $\langle \mathcal{O}_K \rangle$. The dependence on $|\Omega\rangle$ is carried by the functions $f(p)$ and $h^{(l)}(p)$.

We found it more convenient to work with the dimensionful formula (6.7), because this way non-trivial checks of dimensional analysis can be performed at each step of the calculation. The dimensionless formulas can be obtained as explained in 7.5.

The main idea behind the factorization procedure is simple: at each step the number of the integrals can be reduced by one using the integral equation (6.5), whenever the prefactors are such that the corresponding variable is present only in one denominator

$$\frac{1}{(p_j - p_k)^2 + c^2} = \frac{1}{2c} \varphi(p_j - p_k).$$

Except from the case $K = 2$ this is not the case, instead the prefactors have to be divided into several terms, in each of which one of the integrals can be performed. This is a non-trivial task with growing complexity as K increases. In the following we present a case-by-case study up until $K = 4$.

¹The case $K = 1$ is trivial and it simply yields the particle density as it should.

7.1 $K = 2$

One has

$$\langle \mathcal{O}_2 \rangle = 4 \left[\int d\tilde{p}_1 d\tilde{p}_2 \frac{p_2}{(p_1 - p_2)^2 + c^2} h^{(0)}(p_1) h^{(1)}(p_2) - \int d\tilde{p}_1 d\tilde{p}_2 \frac{p_1}{(p_1 - p_2)^2 + c^2} h^{(0)}(p_1) h^{(1)}(p_2) \right].$$

In the first term one integrates over p_1 first, in the second term over p_2 first. Using the integral equation (6.5) one gets

$$\begin{aligned} \langle \mathcal{O}_2 \rangle &= \frac{2}{c} \left[\int d\tilde{p}_2 p_2 h^{(1)}(p_2) (h^{(0)}(p_2) - 1) - \int d\tilde{p}_1 p_1 h^{(0)}(p_1) (h^{(1)}(p_1) - p_1) \right] = \\ &= \frac{2}{c} \int d\tilde{p} (p^2 h^{(0)}(p) - p h^{(1)}(p)). \end{aligned} \quad (7.1)$$

This is in agreement with formula (10) of [35]. As it was already explained in [31, 32, 35], in the finite temperature case (7.1) agrees with the result obtained from the Hellmann-Feynmann theorem. We also note that (7.1) can be proven for arbitrary weight function $f(p)$ using the Hellmann-Feynmann for a single state; one has to repeat the arguments of Appendix D of [34].

For future use we define

$$\{n, m\} = \int d\tilde{p} p^n h^{(m)}(p) = \int \frac{dp}{2\pi} f(p) p^n h^{(m)}(p). \quad (7.2)$$

It can be shown using the iterative solution to (6.5) that in general

$$\{n, m\} = \{m, n\}.$$

Using this notation we write

$$\langle \mathcal{O}_2 \rangle = \frac{2}{c} \left(\{0, 2\} - \{1, 1\} \right). \quad (7.3)$$

Also, it is useful to derive a general formula which will be used often:

$$\int d\tilde{x} d\tilde{y} \frac{x^\alpha - y^\alpha}{(x - y)^2 + c^2} (h^{(\beta)}(x) h^{(\gamma)}(y) - h^{(\beta)}(y) h^{(\gamma)}(x)) = \frac{1}{c} \left(\{\gamma, \alpha + \beta\} - \{\beta, \alpha + \gamma\} \right). \quad (7.4)$$

7.2 $K = 3$

The mean value is given by

$$\langle \mathcal{O}_3 \rangle = 36 \int d\tilde{p}_1 \dots d\tilde{p}_3 \prod_{j>l} \frac{p_j - p_l}{(p_j - p_l)^2 + c^2} h^{(0)}(p_1) h^{(1)}(p_2) h^{(2)}(p_3). \quad (7.5)$$

In this form neither of the integrals can be done directly. Instead, one has to divide the prefactors into several terms such that in each of them one integral can be performed. One way to do this is as follows. We define

$$D(x, y, z) = \frac{y - x}{(x - y)^2 + c^2} \frac{1}{(y - z)^2 + c^2}.$$

Then we find the identity

$$\begin{aligned} \prod_{j>l} \frac{p_j - p_l}{(p_j - p_l)^2 + c^2} &= \\ \frac{1}{3} [D(x, y, z) + D(y, z, x) + D(z, x, y) - D(y, x, z) - D(x, z, y) - D(z, y, x)]. \end{aligned} \quad (7.6)$$

For simplicity we used the variables x, y, z on the r.h.s. instead of p_1, p_2, p_3 . Equation (7.6) can be proven as follows. The r.h.s. is a completely anti-symmetric function of the variables

x, y, z and it has exactly the same poles as the function on the l.h.s., therefore it has to be equal to the l.h.s. multiplied by a symmetric polynomial. By power counting it is shown that this polynomial is a pure number. This number is found to be 1 by simple manipulations after sending $c \rightarrow 0$ on both sides.

Substituting (7.6) into (7.5), performing one integral in each term using (6.5), changing variables accordingly and observing the cancellation of the terms including three h functions we find

$$\begin{aligned} \langle \mathcal{O}_3 \rangle = \frac{6}{c} \int d\tilde{x} d\tilde{y} \frac{y-x}{(x-y)^2 + c^2} \times [y^2(h_1(x)h_0(y) - h_0(x)h_1(y)) + \\ y(h_0(x)h_2(y) - h_2(x)h_0(y)) + \\ (h_2(x)h_1(y) - h_1(x)h_2(y))]. \end{aligned} \quad (7.7)$$

Using (7.4) the last line of (7.7) yields

$$\frac{6}{c} \int d\tilde{x} d\tilde{y} \frac{y-x}{(x-y)^2 + c^2} (h_2(x)h_1(y) - h_1(x)h_2(y)) = \frac{6}{c^2} (\{2, 2\} - \{1, 3\}).$$

For the second line of (7.7) we can drop the term proportional to xy to find

$$\frac{6}{c} \int d\tilde{x} d\tilde{y} \frac{y^2}{(x-y)^2 + c^2} (h_0(x)h_2(y) - h_2(x)h_0(y)) = \frac{3}{c^2} (\{0, 4\} - \{2, 2\}).$$

The first line of (7.7) is more involved. We have to compute

$$\frac{3}{c} \int d\tilde{x} d\tilde{y} \frac{(y-x)(y^2+x^2)}{(x-y)^2 + c^2} (h_1(x)h_0(y) - h_0(x)h_1(y)). \quad (7.8)$$

We write

$$\frac{(y-x)(y^2+x^2)}{(x-y)^2 + c^2} = \frac{y-x}{3} + \frac{c^2}{3} \frac{x-y}{(x-y)^2 + c^2} + \frac{2}{3} \frac{y^3-x^3}{(x-y)^2 + c^2}. \quad (7.9)$$

The first term in (7.9) gives

$$\frac{2}{c} (\{0, 1\}^2 - \{0, 0\}\{1, 1\}).$$

The second and third terms in (7.9) can be evaluated using (7.4).

Putting everything together

$$\langle \mathcal{O}_3 \rangle = \frac{1}{c^2} (-4\{1, 3\} + 3\{2, 2\} + \{0, 4\}) + (\{0, 2\} - \{1, 1\}) + \frac{2}{c} (\{0, 1\}^2 - \{0, 0\}\{1, 1\}). \quad (7.10)$$

This is in accordance with formula (11) of [35].

7.3 $K = 4$

We define

$$D_4(x, y, z, u) = \frac{1}{((x-y)^2 + c^2)((y-z)^2 + c^2)((z-u)^2 + c^2)}.$$

Then we find

$$\prod_{j>l} \frac{p_j - p_l}{(p_j - p_l)^2 + c^2} = \frac{1}{12} \sum_{\mathcal{P} \in \sigma_4} (-1)^{[\mathcal{P}]} D_4(\mathcal{P}p). \quad (7.11)$$

This equation can be proven through the same steps as in the case of (7.6).

One has to evaluate

$$\begin{aligned} \langle \mathcal{O}_4 \rangle &= 24^2 \int d\tilde{p}_1 \dots d\tilde{p}_4 \prod_{j<l} \frac{p_j - p_l}{((p_j - p_l)^2 + c^2)} h^{(0)}(p_1) h^{(1)}(p_2) h^{(2)}(p_3) h^{(3)}(p_4) = \\ &= 48 \int d\tilde{p}_1 \dots d\tilde{p}_4 \frac{1}{((p_1 - p_2)^2 + c^2)((p_2 - p_3)^2 + c^2)((p_3 - p_4)^2 + c^2)} \sum_{\mathcal{P} \in \sigma_4} (-1)^{[\mathcal{P}]} \prod_{l=1}^4 h^{(\mathcal{P}_l - 1)}(p_l). \end{aligned}$$

Performing the integrals over p_1 and p_4 and using the symmetries one gets

$$\langle \mathcal{O}_4 \rangle = \frac{12}{c^2} \sum_{\mathcal{P} \in \sigma_4} (-1)^{[\mathcal{P}]} \int d\tilde{p}_2 d\tilde{p}_3 \frac{p_2^{\mathcal{P}_1-1} p_3^{\mathcal{P}_4-1}}{(p_2 - p_3)^2 + c^2} h^{(\mathcal{P}_2-1)}(p_2) h^{(\mathcal{P}_3-1)}(p_3).$$

There are in total 6 different combinations and we treat them one by one. We define

$$\langle \mathcal{O}_4 \rangle = \frac{12}{c^2} \sum_{o=1}^6 K_o.$$

with K_o representing one of the six combinations. The first three cases are evaluated easily:

$$K_1 = \int d\tilde{x} d\tilde{y} \frac{y-x}{(x-y)^2 + c^2} (h^{(2)}(x)h^{(3)}(y) - h^{(3)}(x)h^{(2)}(y)) = \frac{1}{c} (\{2, 4\} - \{3, 3\})$$

$$K_2 = \int d\tilde{x} d\tilde{y} \frac{y^2 - x^2}{(x-y)^2 + c^2} (h^{(3)}(x)h^{(1)}(y) - h^{(1)}(x)h^{(3)}(y)) = \frac{1}{c} (\{3, 3\} - \{1, 5\})$$

$$K_3 = \int d\tilde{x} d\tilde{y} \frac{y^3 - x^3}{(x-y)^2 + c^2} (h^{(1)}(x)h^{(2)}(y) - h^{(2)}(x)h^{(1)}(y)) = \frac{1}{c} (\{1, 5\} - \{2, 4\})$$

The remaining three case are more complicated because we have to separate factors of the form

$$\frac{x^\alpha y^\beta - x^\beta y^\alpha}{(x-y)^2 + c^2} \quad \alpha, \beta > 0.$$

The next case is

$$K_4 = \int d\tilde{x} d\tilde{y} \frac{xy^2 - yx^2}{(x-y)^2 + c^2} (h^{(0)}(x)h^{(3)}(y) - h^{(3)}(x)h^{(0)}(y)).$$

Here we use

$$\frac{xy^2 - x^2y}{(x-y)^2 + c^2} = \frac{x-y}{3} + \frac{c^2}{3} \frac{y-x}{(x-y)^2 + c^2} + \frac{1}{3} \frac{y^3 - x^3}{(x-y)^2 + c^2}$$

leading to

$$K_4 = \frac{2}{3} (\{0, 1\}\{0, 3\} - \{0, 0\}\{1, 3\}) + \frac{c}{3} (\{0, 4\} - \{1, 3\}) + \frac{1}{3c} (\{0, 6\} - \{3, 3\}).$$

The next case is

$$K_5 = \int d\tilde{x} d\tilde{y} \frac{xy^3 - yx^3}{(x-y)^2 + c^2} (h^{(2)}(x)h^{(0)}(y) - h^{(0)}(x)h^{(2)}(y)).$$

Here we use

$$\frac{xy^3 - x^3y}{(x-y)^2 + c^2} = \frac{1}{2} \left[x^2 - y^2 - c^2 \frac{x^2 - y^2}{(x-y)^2 + c^2} + \frac{y^4 - x^4}{(x-y)^2 + c^2} \right]$$

which gives

$$K_5 = (\{2, 2\}\{0, 0\} - \{0, 2\}^2) + \frac{c}{2} (\{2, 2\} - \{0, 4\}) + \frac{1}{2c} (\{2, 4\} - \{0, 6\}).$$

Finally, the last term is

$$K_6 = \int d\tilde{x} d\tilde{y} \frac{x^2y^3 - y^2x^3}{(x-y)^2 + c^2} (h^{(0)}(x)h^{(1)}(y) - h^{(1)}(x)h^{(0)}(y)).$$

Here we write

$$\begin{aligned} \frac{x^2y^3 - x^3y^2}{(x-y)^2 + c^2} = \frac{1}{5} & \left[\frac{2c^2}{3}(x-y) + (x^3 - y^3) + 2(x^2y - y^2x) + \right. \\ & \left. + \frac{2c^4}{3} \frac{y-x}{(x-y)^2 + c^2} + \frac{5c^2}{3} \frac{y^3 - x^3}{(x-y)^2 + c^2} + \frac{y^5 - x^5}{(x-y)^2 + c^2} \right] \end{aligned}$$

leading to

$$\begin{aligned} K_6 = & \frac{4c^2}{15} \left(\{0, 1\}^2 - \{0, 0\}\{1, 1\} \right) + \frac{2}{5} \left(\{0, 3\}\{0, 1\} - \{1, 3\}\{0, 0\} \right) + \\ & \frac{4}{5} \left(\{0, 2\}\{1, 1\} - \{0, 1\}\{1, 2\} \right) + \frac{2c^3}{15} \left(\{0, 2\} - \{1, 1\} \right) + \frac{c}{3} \left(\{0, 4\} - \{1, 3\} \right) + \\ & \frac{1}{5c} \left(\{0, 6\} - \{1, 5\} \right). \end{aligned}$$

Putting everything together

$$\begin{aligned} \langle \mathcal{O}_4 \rangle = & \frac{2}{5c^3} \left[8c^3 \left(\{0, 1\}^2 - \{0, 0\}\{1, 1\} \right) + 32c \left(\{0, 1\}\{0, 3\} - \{0, 0\}\{1, 3\} \right) + \right. \\ & 24c \left(\{0, 2\}\{1, 1\} - \{0, 1\}\{1, 2\} \right) + 30c \left(\{0, 0\}\{2, 2\} - \{0, 2\}^2 \right) + \\ & 4c^4 \left(\{0, 2\} - \{1, 1\} \right) + 5c^2 \left(\{0, 4\} - 4\{1, 3\} + 3\{2, 2\} \right) + \\ & \left. \{0, 6\} - 6\{1, 5\} + 15\{2, 4\} - 10\{3, 3\} \right]. \end{aligned} \quad (7.12)$$

This a new result of the present work.

7.4 Galilei invariance

The expectation values $\langle \mathcal{O}_K \rangle$ are Galilei invariant, and it is useful to check this property in our final formulas. This constitutes a highly non-trivial check, as it was already remarked in [35].

In our calculations we did not restrict ourselves to symmetric distributions, the weight functions $f(p)$ can be arbitrary. Therefore, to check Galilei invariance it is enough to consider an infinitesimal boost b . This boost yields the following infinitesimal transformations:

$$p \rightarrow p + b \quad h^{(j)}(p) \rightarrow h^{(j)}(p) + bj h^{(j-1)}(p).$$

It is then readily seen that (6.7) is invariant due to the anti-symmetry of the prefactors.

We also performed the check on our factorized formulas. The transformation rules for the quantities $\{\alpha, \beta\}$ are

$$\{\alpha, \beta\} \rightarrow \{\alpha, \beta\} + b \left[\alpha \{\alpha - 1, \beta\} + \beta \{\alpha, \beta - 1\} \right].$$

Using this rule we have checked that the variation of equations (7.3), (7.10) and (7.12) indeed vanishes.

7.5 Dimensionless formulas and numerical results

The dimensionless versions of formulas (7.3), (7.10) and (7.12) are obtained simply by setting $c = 1$, multiplying with an overall factor of γ^K , and replacing

$$\{n, m\} \rightarrow \int d\tilde{q} \, q^n \tilde{h}^{(m)}(q), \quad (7.13)$$

To numerically evaluate the factorized formulas the following steps have to be performed:




Figure 1: The ground state values of the K -body local correlations for $K \leq 4$ as a function of the dimensionless coupling γ ($g_1 = 1$ by definition). The exact values are represented by the solid lines, whereas the dashed lines show the empirical formula (7.14).

- Solve the TBA equation (6.8) iteratively. The parameter α can be fixed by requiring

$$g_1 = \gamma\{0, 0\} = 1$$

- Solve the linear integral equations (6.9) for $\tilde{h}^{(l)}(q)$.
- Evaluate (7.3), (7.10) and (7.12).

We performed this procedure for a wide range of the parameters γ and τ . The quantity g_4 shows the same qualitative behaviour as g_2 and g_3 [35]: it is an increasing function of τ and a decreasing function of γ , with the limiting values given by

$$\lim_{\gamma \rightarrow 0} g_4 = \lim_{\tau \rightarrow \infty} g_4 = 4! = 24 \quad \lim_{\gamma \rightarrow 0} \lim_{\tau \rightarrow 0} g_4 = 1.$$

To demonstrate the numerical results we present the ground state values of g_2 , g_3 and g_4 in Fig. 1, whereas the temperature dependence of g_4 is shown in Fig. 2 for the intermediate couplings $\gamma = 0.1$, $\gamma = 1$ and $\gamma = 10$.

At $T = 0$ the first term in the small coupling expansion of g_K is given by [29]:

$$g_K = 1 - \frac{K(K-1)}{\pi} \sqrt{\gamma} + \mathcal{O}(\gamma)$$

We found that the empirical formula

$$g_K \approx \exp\left(-\frac{K(K-1)}{\pi} \sqrt{\gamma}\right) \quad (7.14)$$

gives a surprisingly good approximation and can be used for practical purposes even at $\gamma \sim 1$. The predictions of (7.14) are also plotted in Fig. 1. It is expected that (7.14) holds with a good approximation even for higher K .

It would be useful to compare the exact numerical values for g_3 and g_4 case to the various approximations available in the literature [28, 29] including the large-coupling expansion both at zero and finite temperatures. This is out of the scope of the present work and is left for further research.




Figure 2: The quantity g_4 as a function of the dimensionless temperature τ for intermediate couplings. In the $\tau \rightarrow 0$ limit the three curves approach small (but non-zero) values which are shown in Figure 1.

8 Mean values in the LeClair-Mussardo formalism

In this section we elaborate on the LeClair-Mussardo formalism, which is an alternative approach to obtain expectation values of local operators leading to an infinite integral series [51, 52, 31, 32, 34]. For our present purposes the following form of the series is the most convenient [34]:

$$\langle \mathcal{O}_K \rangle = \sum_N \frac{1}{N!} \int \frac{dp_1}{2\pi} \cdots \frac{dp_N}{2\pi} \left(\prod_{j=1}^N f(p_j) \omega(p_j) \right) F_{N,s}^K(p_1, \dots, p_N), \quad (8.1)$$

where

$$\omega(p) = \exp \left(- \int \frac{dp'}{2\pi} f(p') \varphi(p - p') \right)$$

and $f(p)$ is defined in (6.2). The form factors appearing in the above series are defined as

$$F_{N,s}^K(p_1, \dots, p_N) = \prod_{j < k} \frac{(p_j - p_k)^2}{(p_j - p_k)^2 + c^2} \times \lim_{\varepsilon \rightarrow 0} F_N^K(\{p_j + \varepsilon\}, \{p_j\}), \quad (8.2)$$

where the form factor on the r.h.s. is given by (5.4). This prescription is also called the “symmetric evaluation of the diagonal limit”; note that this limit is different from the way we obtained the mean value (5.6) because in (5.4) the Bethe equations were substituted into the matrix element *before* taking the diagonal limit. Therefore the object in (8.2) does not depend on the volume L . Note also that the l.h.s. refers to a normalization where the norm of the Bethe state is given simply by the Gaudin-determinant $\mathcal{G}^{LL}$.

Alternatively (8.1) can be expressed as [34, 31, 32]

$$\langle \mathcal{O}_K \rangle = \sum_N \frac{1}{N!} \int \frac{dp_1}{2\pi} \cdots \frac{dp_N}{2\pi} \left(\prod_{j=1}^N f(p_j) \right) F_{N,c}^K(p_1, \dots, p_N). \quad (8.3)$$

Here $F_{N,c}^K(p_1, \dots, p_N)$ are the so-called connected evaluations of the diagonal form factors. Their precise definition and the relation to $F_{N,s}^K(p_1, \dots, p_N)$ can be found in [52, 34]. The series (8.3) was originally developed in [51] in the framework of integrable Quantum Field Theories. Later it was used in [31, 32, 35] to compute the quantities g_K up to $K = 3$.

However, for higher K the results (8.1)-(8.3) are only formal because the form factors themselves were not calculated previously. In [31, 32] a prescription was given of how to obtain the connected evaluation using a special non-relativistic limit of certain form factors of the sinh-Gordon model. However, the actual calculation becomes more and more demanding with higher K and N .

We fill this gap here by calculating the explicit results for $F_{N,s}^K(p_1, \dots, p_N)$ for arbitrary K and N : we take the symmetric diagonal limit of the form factor (5.4). Note that every singularity of the form factor is included in the matrix Z , and even the elements of Z can be evaluating easily. Taking the limit and multiplying with the prefactors we obtain

$$F_{N,s}^K(p_1, \dots, p_m) = (K!)^2 \sum_{\substack{\{p^+\} \cup \{p^-\} \\ |\{p^+\}|=K}} \left[\prod_{j>l} \frac{p_j^+ - p_l^+}{(p_j^+ - p_l^+)^2 + c^2} \right] \times \det Y. \quad (8.4)$$

The elements of Y are given by

$$Y_{jl} = (p_j)^{l-1} \quad \text{if } p_l \in \{p^+\}$$

$$Y_{j,l} = \delta_{j,l} \left(\sum_{o=1}^N \varphi(p_j - p_o) \right) - \varphi(p_j - p_l) \quad \text{if } p_l \in \{p^-\}. \quad (8.5)$$

Note that

$$Y = \mathcal{H} \Big|_{L=0},$$

where $\mathcal{H}$ is the matrix defined in (5.7).

With these results the series (8.1) can be considered an explicit representation of the mean value.

It would be desirable to have a general recipe for the re-summation of the series, which would be an alternative way to obtain factorized formulas like (7.10) and (7.12). However, this is far from being easy. The simpler cases $K = 1$ and $K = 2$ were already calculated in [31, 32]. The highly non-trivial case of $K = 3$ was considered in [35], where the authors evaluated the series (8.3) (and obtained the result (7.10) for the first time) based on the following conjecture for the quantities $F_{N,c}^3$ ²:

$$F_{N,c}^3 = \frac{1}{2c^2} \sum_P \varphi_{12} \varphi_{23} \dots \varphi_{N-1,N} p_{1N} (p_{1N}^3 - p_{12}^3 - p_{23}^3 - \dots - p_{N-1,N}^3). \quad (8.6)$$

Here we check this formula in the first two cases. In the simplest case of $N = 3$ our formula (8.4) gives

$$F_{3,s}^3 = F_{3,c}^3 = 36 \prod_{j>l} \frac{p_{jl}^2}{p_{jl}^2 + c^2}.$$

This was already calculated in [31, 32] and is in agreement with (8.6). In the case of $N = 4$ one has to use the following relation between the symmetric and connected evaluations [52]:

$$F_{4,s}^3(p_1, p_2, p_3, p_4) = F_{4,c}^3(p_1, p_2, p_3, p_4) + \sum_j F_{3,c}^3(\hat{p}_j) \times \left(\sum_{k \neq j} \varphi_{jk} \right).$$

Here $\hat{p}_j$ means that p_j is not present among the arguments of the form factor. We used the program **Mathematica** to express $F_{4,c}^3$ using the above relation and we found agreement with (8.6). This is a highly non-trivial check of the conjecture (8.6); a proof for arbitrary N is not known.

Finally we note that in the simpler cases of $K = 1$ and $K = 2$ we evaluated (8.4) and found exact agreement with the corresponding formulas of Appendix D in [34].

²In order to ensure compatibility with our normalizations we inserted a factor of $1/c^2$.

9 Conclusions

We developed multiple integral formulas for the local correlations in the 1D Bose gas. The final results for the expectation value $\langle \mathcal{O}_K \rangle$ is given by equation (6.7), whereas the dimensionless formula for g_K is given by (6.10).

In section 7 we performed the explicit factorization of the multiple integrals in the cases $K = 2, 3, 4$; for $K = 3$ we obtained the recent result of [35] whereas our formula for $K = 4$ is new. Our method of factorization relies only on the integral equation (6.5) defining the auxiliary functions entering the multiple integral. Therefore the process works for arbitrary distribution of Bethe roots and not only for the ground state or the finite temperature Gibbs states.

The general recipe of how to perform the factorization for $K > 4$ is not known. The strategy is clear: at each step the prefactors have to be manipulated in such a way that the number of integrals can be reduced by one using the integral equation (6.5). We believe that this can always be done and it would be interesting to develop a general algorithm for this process.

An alternative way to obtain the mean values $\langle \mathcal{O}_K \rangle$ would be to take the thermodynamic limit on the XXZ spin chain first and to perform the scaling limit towards the Bose gas afterwards. The advantage of this approach would be that on the spin chain the factorization of the multiple integral formulas for the elements of the reduced density matrix is by now well-understood (see [53] and references therein). In fact we attempted to take the scaling limit of the factorized results of [54] concerning the emptiness formation probability. However, this turned out to be cumbersome already in the case $K = 2$. Thus it seems that the direct approach of the present paper is more advantageous, at least for the small values of K considered here.

An other alternative way would be to sum up the LeClair-Mussardo series (8.1) or (8.3). The diagonal form factors entering (8.1) are given explicitly by (8.4), therefore the remaining task is purely combinatorial: one has to expand the sums of determinants appearing in (8.4) and put the resulting expression in a form which is amenable for re-summation. Again, this is a formidable problem, the solution of which is not yet known.

Acknowledgements The author is grateful to M. Kormos and A. Imambekov for inspiring discussions and for communicating the results of [35] prior to publication. Also, we are indebted to J. Mossel, N. J. van Druten, J-S. Caux and M. Kormos for useful comments about the manuscript.

References

- [1] E. H. Lieb and W. Liniger, “Exact Analysis of an Interacting Bose Gas. I. The General Solution and the Ground State,” *Phys. Rev.* **130** (1963) 1605.
- [2] E. H. Lieb, “Exact Analysis of an Interacting Bose Gas. II. The Excitation Spectrum,” *Phys. Rev.* **130** (1963) no. 4, 1616–1624.
- [3] H. Bethe, “Zur Theorie der Metalle,” *Zeitschrift für Physik* **A71** (1931) 205.
- [4] C. N. Yang and C. P. Yang, “Thermodynamics of a One-Dimensional System of Bosons with Repulsive Delta-Function Interaction,” *J. Math. Phys.* **10** (1969) 1115.
- [5] D. B. Creamer, H. B. Thacker, and D. Wilkinson, “Some exact results for the two-point function of an integrable quantum field theory,” *Phys. Rev. D* **23** (1981) no. 12, 3081–3084.
- [6] D. B. Creamer, H. B. Thacker, and D. Wilkinson, “A study of correlation functions for the delta-function Bose gas,” *Physica D* **20** (1986) 155.
- [7] V. Korepin, N. Bogoliubov, and A. Izergin, *Quantum inverse scattering method and correlation functions*. Cambridge University Press, 1993.

- [8] N. A. Slavnov, “Nonequal-time current correlation function in a one-dimensional Bose gas,” *Theoretical and Mathematical Physics* **82** (1990) 273–282.
- [9] N. Kitanine, K. Kozłowski, J. M. Maillet, N. A. Slavnov, and V. Terras, “On correlation functions of integrable models associated with the six-vertex R-matrix,” *J. Stat. Mech.* (2007) P01022, [arXiv:hep-th/0611142](#).
- [10] J. Caux and P. Calabrese, “Dynamical density-density correlations in the one-dimensional Bose gas,” *Phys. Rev. A* **74** (2006) no. 3, 031605, [arXiv:cond-mat/0603654](#).
- [11] A. Seel, T. Bhattacharyya, F. Göhmann, and A. Klümper, “A note on the spin-1/2 XXZ chain concerning its relation to the Bose gas,” *J. Stat. Mech.* (2007) P08030, [arxiv:0705.3569](#).
- [12] K. K. Kozłowski, J. M. Maillet, and N. A. Slavnov, “Long-distance behavior of temperature correlation functions in the one-dimensional Bose gas,” *J. Stat. Mech.* **3** (2011) 18, [arXiv:1011.3149 \[math-ph\]](#).
- [13] K. K. Kozłowski, J. M. Maillet, and N. A. Slavnov, “Correlation functions for one-dimensional bosons at low temperature,” *J. Stat. Mech.* **3** (2011) 19, [arXiv:1102.1858 \[math-ph\]](#).
- [14] A. Imambekov and L. I. Glazman, “Exact Exponents of Edge Singularities in Dynamic Correlation Functions of 1D Bose Gas,” *Phys. Rev. Lett.* **100** (2008) no. 20, 206805, [arXiv:0711.2697 \[cond-mat.str-el\]](#).
- [15] A. Shashi, L. I. Glazman, J.-S. Caux, and A. Imambekov, “Exact asymptotes of static and dynamic correlation functions of the 1D Bose gas,” [arXiv:1010.2268 \[cond-mat.stat-mech\]](#).
- [16] T. Kinoshita, T. Wenger, and D. S. Weiss, “Observation of a One-Dimensional Tonks-Girardeau Gas,” *Science* **305** (2004) no. 5687, 1125–1128.
- [17] T. Kinoshita, T. Wenger, and D. S. Weiss, “A quantum Newton’s cradle,” *Nature* **440** (2006) 900.
- [18] D. Petrov, D. Gangardt, and G. Shlyapnikov, “Low-dimensional trapped gases,” *J. Phys. IV France* **116** (2004) 5, [arXiv:cond-mat/0409230](#).
- [19] A. H. van Amerongen, J. J. P. van Es, P. Wicke, K. V. Kheruntsyan, and N. J. van Druten, “Yang-Yang Thermodynamics on an Atom Chip,” *Phys. Rev. Lett.* **100** (2008) no. 9, 090402, [arXiv:0709.1899 \[cond-mat.other\]](#).
- [20] S. Trotzky, Y.-A. Chen, A. Flesch, I. P. McCulloch, U. Schollwöck, J. Eisert, and I. Bloch, “Probing the relaxation towards equilibrium in an isolated strongly correlated 1D Bose gas,” [arXiv:1101.2659 \[cond-mat.quant-gas\]](#).
- [21] V. Guarrera, P. Wuertz, A. Ewerbeck, A. Vogler, G. Barontini, and H. Ott, “Observation of local temporal correlations in trapped quantum gases,” [arXiv:1105.4818](#).
- [22] E. Haller, M. Rabie, M. J. Mark, J. G. Danzl, R. Hart, K. Lauber, G. Pupillo, and H.-C. Nägerl, “Three-body correlation functions and recombination rates for bosons in three and one dimensions,” [arXiv:1107.4516 \[cond-mat.quant-gas\]](#).
- [23] M. J. Davis, P. B. Blakie, A. H. van Amerongen, N. J. van Druten, and K. V. Kheruntsyan, “Yang-Yang thermometry and momentum distribution of a trapped one-dimensional Bose gas,” [arXiv:1108.3608 \[cond-mat.quant-gas\]](#).
- [24] Y. Kagan, B. V. Svistunov, and G. V. Shlyapnikov, “Effect of Bose condensation on inelastic processes in gases,” *JETP letters* **42** (1985) 209.
- [25] B. L. Tolra, K. M. O’Hara, J. H. Huckans, W. D. Phillips, S. L. Rolston, and J. V. Porto, “Observation of Reduced Three-Body Recombination in a Correlated 1D Degenerate Bose Gas,” *Phys. Rev. Lett.* **92** (2004) 190401.

- [26] J. Armijo, T. Jacqmin, K. V. Kheruntsyan, and I. Bouchoule, “Probing Three-Body Correlations in a Quantum Gas Using the Measurement of the Third Moment of Density Fluctuations,” *Phys. Rev. Lett.* **105** (2010) 230402, [arXiv:1007.3713](#).
- [27] T. Jacqmin, J. Armijo, T. Berrada, K. V. Kheruntsyan, and I. Bouchoule, “Sub-Poissonian Fluctuations in a 1D Bose Gas: From the Quantum Quasicondensate to the Strongly Interacting Regime,” *Phys. Rev. Lett.* **106** (2011) 230405, [arXiv:1103.3028](#).
- [28] D. M. Gangardt and G. V. Shlyapnikov, “Stability and phase coherence of trapped 1D Bose gases,” *Phys. Rev. Lett.* **90** (2002) , [arXiv:cond-mat/0207338](#).
- [29] D. M. Gangardt and G. V. Shlyapnikov, “Local correlations in a strongly interacting 1D Bose gas,” *New Journal of Physics* **5** (2003) 79, [arXiv:cond-mat/0304159](#).
- [30] V. V. Cheianov, H. Smith, and M. B. Zvonarev, “Three-body local correlation function in the Lieb Liniger model: bosonization approach,” *J. Stat. Mech.* **8** (2006) 15, [arXiv:cond-mat/0602468](#).
- [31] M. Kormos, G. Mussardo, and A. Trombettoni, “Expectation Values in the Lieb-Liniger Bose Gas,” *Phys. Rev. Lett.* **103** (2009) 210404, [arXiv:0909.1336 \[cond-mat.stat-mech\]](#).
- [32] M. Kormos, G. Mussardo, and A. Trombettoni, “1D Lieb-Liniger Bose Gas as Non-Relativistic Limit of the Sinh-Gordon Model,” *Phys. Rev. A* **81** (2010) 043606, [arXiv:0912.3502 \[cond-mat.stat-mech\]](#).
- [33] M. Kormos, G. Mussardo, and B. Pozsgay, “Bethe Ansatz Matrix Elements as Non-Relativistic Limits of Form Factors of Quantum Field Theory,” *J. Stat. Mech.* (2010) P05014, [arXiv:1002.3387 \[cond-mat.stat-mech\]](#).
- [34] B. Pozsgay, “Mean values of local operators in highly excited Bethe states,” *J. Stat. Mech.* **2011** (2011) P01011, [arXiv:1009.4662 \[hep-th\]](#).
- [35] M. Kormos, Y.-Z. . Z. Chou, and A. Imambekov, “Exact three-body local correlations for arbitrary excited states of the 1D Bose gas,” [arXiv:1108.0145](#).
- [36] B. Golzer and A. Holz, “The nonlinear Schrodinger model as a special continuum limit of the anisotropic Heisenberg model,” *J. Phys. A* **20** (1987) no. 11, 3327.
- [37] N. Kitanine, J. M. Maillet, N. A. Slavnov, and V. Terras, “Spin-spin correlation functions of the XXZ-1/2 Heisenberg chain in a magnetic field,” *Nucl. Phys. B* **641** (2002) 487–518, [arXiv:hep-th/0201045](#).
- [38] N. Kitanine, J. M. Maillet, and V. Terras, “Form factors of the XXZ Heisenberg spin-1/2 finite chain,” *Nucl. Phys. B* **554** (1999) 647–678, [arXiv:math-ph/9807020](#).
- [39] M. Gaudin, “La fonction d’onde de Bethe pour les modèles exacts de la mécanique statistique,” *Commisariat à l’énergie atomique, Paris* (1983) .
- [40] V. E. Korepin, “Calculation of norms of Bethe wave functions,” *Comm. Math. Phys.* **86** (1982) 391.
- [41] R. Orbach, “Linear Antiferromagnetic Chain with Anisotropic Coupling,” *Phys. Rev.* **112** (1958) no. 2, 309–316.
- [42] L. R. Walker, “Antiferromagnetic Linear Chain,” *Phys. Rev.* **116** (1959) no. 5, 1089–1090.
- [43] C. N. Yang and C. P. Yang, “One-Dimensional Chain of Anisotropic Spin-Spin Interactions. I. Proof of Bethe’s Hypothesis for Ground State in a Finite System,” *Phys. Rev.* **150** (1966) no. 1, 321–327.
- [44] J. H. H. Perk, “Bethe-Ansatz en Eendimensionale Quantumsystemen,” Master’s thesis, University of Amsterdam, 1974.
- [45] F. Göhmann and V. E. Korepin, “Solution of the quantum inverse problem,” *J. Phys. A* **33** (2000) no. 6, 1199, [arXiv:hep-th/9910253](#).

- [46] N. A. Slavnov, “Calculation of scalar products of wave functions and form factors in the framework of the algebraic Bethe ansatz,” *Theoretical and Mathematical Physics* **79** (1989) 502–508.
- [47] A. G. Izergin, D. A. Coker, and V. E. Korepin, “Determinant formula for the six-vertex model,” *J. Phys. A* **25** (1992) no. 16, 4315.
- [48] J. Mossel and J. Caux, “Relaxation dynamics in the gapped XXZ spin-1/2 chain,” *New Journal of Physics* **12** (2010) no. 5, 055028, [arXiv:1002.3988 \[cond-mat.str-el\]](#).
- [49] V. E. Korepin and N. A. Slavnov, “Form Factors in the Finite Volume,” *Int. J. Mod. Phys. B* **13** (1999) 2933–2941, [arXiv:math-ph/9812026](#).
- [50] N. Kitanine, J. M. Maillet, and V. Terras, “Correlation functions of the XXZ Heisenberg spin-1/2 chain in a magnetic field,” *Nucl. Phys. B* **567** (2000) no. 3, 554 – 582, [arXiv:math-ph/9907019](#).
- [51] A. Leclair and G. Mussardo, “Finite temperature correlation functions in integrable QFT,” *Nucl. Phys.* **B552** (1999) 624–642, [arXiv:hep-th/9902075](#).
- [52] B. Pozsgay and G. Takacs, “Form factors in finite volume II: disconnected terms and finite temperature correlators,” *Nucl. Phys.* **B788** (2008) 209–251, [arXiv:0706.3605 \[hep-th\]](#).
- [53] J. Sato, B. Aufgebauer, H. Boos, F. Göhmann, A. Klümper, M. Takahashi, and C. Trippe, “Computation of static Heisenberg-chain correlators: Control over length and temperature dependence,” (2011) , [arXiv:1105.4447](#).
- [54] H. E. Boos, F. Göhmann, A. Klümper, and J. Suzuki, “Factorization of the finite temperature correlation functions of the XXZ chain in a magnetic field,” *J. Phys. A* **40** (2007) 10699, [arXiv:0705.2716](#).